

Lathyrus Production Technology



सत्यमेव जयते

Government of India

Ministry of Agriculture & Farmers Welfare

Department of Agriculture Cooperation & Farmers Welfare

Directorate of Pulses Development, Bhopal (M.P.)



एक कदम स्वच्छता की ओर



Per Drop, More Crop

M-kisan portal - <http://mkisan.gov.in>

Farmers portal - <http://farmer.gov.in>

Kisan Call Centre (KCC)-Toll Free No.-1800-180-1551

LATHYRUS



- Botanical Name - *Lathyrus sativus* (L.)
Synonym - Grass pea, Chickling pea,
Khesari, Teora, Kasari (bengali)
Origin - South Europe and Western Asia

Importance

Lathyrus is considered as drought-tolerant hardy crop, and is grown in low-rainfall regions under rainfed conditions, during winter when lentil and chickpea are not expected to give good yields. The crop has unique tolerance ability against stress environmental conditions not only drought but also for water logging. In addition to use as dal and chapatti, it is usually grown as fodder crop. Lathyrus leaves about 36-48 kg/ha nitrogen economy for the succeeding crop.

Nutritive Value

Protein	-	31.9%	Fat	-	0.9%
Carbohydrate	-	53.9%	Ash	-	3.2%

Crop Status

During Twelfth Plan (2012-2015) the total area and production of Khesari were recorded at 4.93 lakh hectares and 3.84 lakh tonnes respectively. Chattisgarh ranked first both in area and production (67.26 % and 59.52%), followed by Bihar (13.62 % and 20.09%). Madhya Pradesh ranked third in area (8.80%), whereas in production W.B. ranked IIIrd (9.56%), due to highest yield among the all lathyrus producing state.

Climate Requirement

Being a winter season crop it prefers temperate climate with good adoption under climatic extremities. Generally, crop requires 15 °C to 25 °C temperatures during sowing to harvesting of crop.

Varieties : Bio L-212 (Ratan), Prateek, Maha Teora,

Soil Type and Field Preparation

Thrives well in all types of soils except high acidic soils. It prefers heavy soils belonging to low lying areas which are not suited to other crops. It grows abundantly in loamy and deep black soils and produce

excellent crop. For cultivation of lathyrus under *utera* system (relay cropping), no tillage is required. However, for planting after harvest of rice, one deep ploughing followed by cross harrowing and planking is necessary.

Cropping System

It is grown as single crop of the year in areas where water gets accumulated during rainy season or as a relay crop after paddy often as *utera* / *paira* crop in standing paddy, due to its ability to withstand in high moisture conditions at sowing time and moisture stress during growth period.

Sowing Time

Crop is sown on residual soil moisture after harvest of kharif during last October to early November as pure crop. In *utera* cropping last week of September or first week of October.

Seed Rate & Spacing

70-80 kg/ha for broadcasted sowing in *utera* system and 40-60 kg/ha in line sowing is required. Under *utera* cropping sown as broadcasted in-between the rice rows. Whereas normal spacing 30 cm x 10 cm is recommended.

Seed Treatment

Treat the seed with Thiram 3g / kg of seed. After seed treatment with fungicide seed should be inoculated with *Rhizobium* and PSB culture @ 5-7 g/kg of seed.

Plant Nutrient Management

Under *utera* cropping the crop is grown on residual fertility of rice. However, it responds well to phosphorus up to 40-60 kg /ha except in the case if grown on highly phosphorus fertilized paddy field. For normal crop 100 kg DAP + 100 kg gypsum/ha is a optimum dose of fertilizer applied as basal dose 2-3 cm below the seed with the help of ferti-seed drill, is recommended.

Water Management

The crop is grown as rain fed crop on residual moisture. However,

under high moisture stresses one irrigation at 60-70 days after sowing may be remunerative in terms of production.

Weed Management

For normal sown crops one hand-weeding at 30-35 days after sowing (if soil condition permit). Weeds can also be managed effectively by spray of fluchloralin (Basalin) 50% EC @ 0.75-1 kg a.i./ha in 750-1000 liters of water as pre-plant incorporation.

Plant Protection Measures

Insect Pest/Disease/CO	Nature of Damage/ Symptoms	Control Measures
Aphid	The adults and nymphs suck the juice from the leaves as a result, leaves turn brown and crumpled and the plant look sick.	Spray of Dimethoate 30 EC @ 1.7 ml/liter or oxydemeton methy (Metasystox) 25 EC @ 1 ml/liter of water.
Rust (<i>Uromyces fabae</i>)	Pink to brown pustules appeared on leaves and stems. In severe attack, the affected plants are dry.	<ul style="list-style-type: none"> • Grow early maturing variety. • Seed Treatment with Carbendazim @ 2 g/kg seed. • Spray the crop with Mancozeb 75 WP @ 2 g / liter of water.
Downy Mildew (<i>Peronospora spp.</i>)	Brownish cottony growth of fungus may be seen on the lower surface of leaf. Inside growth yellow to greenish spots are also visible.	Spray the crop with Mancozeb 75 WP @ 2 g / liter of water.
Powdery Mildew (<i>Erysiphe polygoni</i>)	Symptoms first appeared on all the aerial part of plant. While powdery masses of spores formed on leaves which may collapse and cover the whole leaf with powdery growth.	Wettable Sulphur @ 3 g/ litre or Carbendazim @ 1g/ liter or Dinocap @ 1 ml/litre of water.

Harvesting, Threshing & Storage

Harvest the crop, when colour of pods change to brown and grains are at dough stage having approximately 15% moisture in-side them. Harvested produce may be allowed to dry in sunlight for a week. Harvested produce after 3-4 days sun drying is roaped in the bundles and transferred to threshing floors. Threshing is done by beating with sticks or trampling under the feet of bullocks. The clean seed should be sun dried for 3-4 days to reduce their moisture content up to 9-10%. Now the produce should be safely stored in appropriate bins. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc).

Yield

A well managed crop can easily give 8-10 q/ha yields under direct sowing and 3-4 q/ha under utera cultivation.

Recommendation to achieved higher production

- i) Deep summer ploughing once in 3 years.
 - ii) Seed treatment should be done before sowing.
 - iii) Application of fertilizer should be based on soil test value.
 - iv) In Lathyrus foliar spray of 2% urea or 20 ppm Salicylic acid at flowering and pod formation stage increases the yield.
 - v) Weed control should be done at right time.
 - vi) Adopt integrated approach for plant protection.
-
- For technical information of crop production please contact to district KVK/ nearest KVK.
 - To avail benefit from Central and State Government running schemes for crop production (ploughing, fertilizers, micronutrient, pesticide, irrigation equipment), agricultural implements, storage infrastructure etc., please contact to your DDA/SADO office.

**- For more information also visit -
M-kisan portal - <http://mkisan.gov.in>
Farmers portal - <http://farmer.gov.in>
Kisan Call Centre (KCC)-Toll Free
No. - 1800-180-1551**

Compiled and edited by

Dr. A.K. Tiwari
Dr. A.K. Shivhare
Shri Vipin Kumar

Technical Support

Shri Sarju Pallearwar
Smt. Ashwini Bhoware

Publisher

Director
Directorate of Pulses Development
Govt. of India
Department of Agriculture
Cooperation & Farmers Welfare
Ministry of Agriculture & Farmers Welfare
6th Floor, Vindhyachal Bhavan
Bhopal – 462004 (M.P.)
E-mail : dpd.mp@nic.in
Telefax : 0755-2571678
Phone : 0755-2550353/ 2572313
Website: www.dpd.gov.in

Publication year 2017